



NAUTILE[®] DG

MAPP No.14692



A water-dispersible granule containing 5% w/w cymoxanil and 68% w/w mancozeb.

For the control of potato blight.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

COMPLIANCE WITH THE FOLLOWING CONDITIONS OF USE AND ALL SAFETY PRECAUTIONS MARKED * IS A LEGAL REQUIREMENT

FOR USE AS AN AGRICULTURAL FUNGICIDE

Crop:	Potato
Maximum Individual Dose:	1.8 kg product / ha
Other specific restrictions:	A minimum of 7 days must be observed between applications

READ ALL OTHER SAFETY PRECAUTIONS AND DIRECTIONS BEFORE USE

SAFETY PRECAUTIONS

Operator Protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

OPERATORS MUST WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate.

However, if engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WASH CONCENTRATE from skin or eyes immediately. **DO NOT BREATHE SPRAY.**

IN CASE OF CONTACT WITH EYES, RINSE IMMEDIATELY with plenty of water and seek medical advice.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.

Environmental Protection

DO NOT CONTAMINATE SURFACE WATERS OR DITCHES with chemical or used container.

Storage and disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

EMPTY CONTAINER COMPLETELY and dispose of safely.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM FIRE AND SPARKS.

A water-dispersible granule containing 5% w/w cymoxanil and 68% w/w mancozeb.

For the control of potato blight.

Xi



IRRITATING TO THE RESPIRATORY SYSTEM. MAY CAUSE SENSITISATION BY SKIN CONTACT.

IRRITANT

Avoid contact with skin.
Wear suitable gloves.
Keep out of reach of children.
Keep away from food, drink and animal feeding stuffs.

N



VERY TOXIC TO AQUATIC ORGANISMS, MAY CAUSE LONG TERM ADVERSE EFFECTS IN THE AQUATIC ENVIRONMENT.

DANGEROUS FOR THE ENVIRONMENT

This material and its container must be disposed of in a safe way.
Use appropriate containment to avoid environmental contamination.

Registration holder and Producer:

Cerexagri B.V.
Tankhoofd 10
3196 KE Vondelingsplaat/Rotterdam
The Netherlands
Tel: +31 10 472 51 00
Fax: +31 10 438 26 13
Email: contact.cerexagri@uniphos.com

Net Contents

10 kg

Nautil[®] is a registered trademark of Cerexagri

Marketing Company:

United Phosphorus Limited

Chadwick House, Birchwood Park, Warrington, WA3 6AE, UK

IN CASE OF TOXIC OR TRANSPORT EMERGENCY PLEASE CALL CARECHEM24 INTERNATIONAL ON 0208 762 8322



DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

NAUTILE® DG is for use on potatoes only.

Disease controlled:

Potato blight (*Phytophthora infestans*)

CROP SPECIFIC INFORMATION:

Rate of use: 1.8 kg product/hectare

Timing:

Spray NAUTILE® DG at first risk of infection. Otherwise, spray as the crop meets within the rows. According to the disease incidence and weather conditions repeat at 7 to 14 days intervals.

The longer interval is recommended under dry conditions. Under the following conditions reduce the spray interval to 10 days:

- On irrigated crops
- On very susceptible varieties e.g. Russet Burbanks
- When active blight is present in areas on crops, dumps or volunteers early in the season
- When the weather favors very rapid disease development

Reduce the spray to 7 days intervals under exceptional circumstances.

Spray until the haulm dies down or is desiccated.

Varieties:

All varieties of early and main crop potatoes may be treated. On crops grown for processing consult your processor before use.

Water volumes:

Not less than 200 litres of water per hectare.

RESISTANCE MANAGEMENT

When fungicides with the same mode of action are used repeatedly over several years in the same field, naturally occurring less sensitive strains may survive, propagate and become dominant in that field. It is therefore advised that wherever possible spray programs should include fungicides with differing modes of action.

FOLLOWING CROPS

Any crop may be sown after a crop treated with NAUTILE® DG.

MIXING

Do not add the chemical directly to the spray tank. Cream the required amount in a small quantity of water, pour and rinse through the filter into a half filled spray tank. Agitate thoroughly whilst filling and throughout the spraying operation.

GUIDANCE RELATING TO PRODUCT USE

- Apply if risk of infection from potato blight, when an official blight warning has been issued or a Blight Infection Period (Smith Period) has occurred.
- Under exceptional circumstances when blight risk is high or rain forecasted during the day following the spray, apply at 7 days intervals.
- Do not keep open pack from one season to another as this may reduce the effectiveness of the fungicide and may create vapors which may be flammable.

All goods supplied are high grade and we believe them to be suitable for the purpose for which we expressly supply them: but as we cannot exercise any control over their mixing, use or application which may affect the performance of the goods all conditions and warranties statutory or otherwise as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us or our Associate Companies for any damage or injury whatsoever arising from their storage, handling, reapplication or use. These conditions cannot be varied by our staff, our agents or the re-sellers of the product whether or not they supervise or assist in the use of such goods